

LA-UR-21-27289

Approved for public release; distribution is unlimited.

Title: Understanding materials in extreme conditions via spectroscopy

Author(s): O'Neal, Kenneth Robert

Intended for: Job interview technical presentation

Issued: 2021-07-26

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.




Understanding materials in extreme conditions via spectroscopy

Ken O'Neal

Spectroscopy

- Study of light-matter interaction
 - Light can be absorbed, transmitted, reflected, refracted, emitted, etc.
- Probes a wide variety of properties depending on wavelength of light
- Can be static or dynamic




Outline

- **FTIR and Raman**
 - Pressure-induced dimensionality crossover in Cu-3Cl(pyd)
 - Local structure changes in Ni_3TeO_6 under pressure
 - Size-dependent vibronic coupling in CuGeO_3 nanorods
- Ultrafast spectroscopy
 - Carrier dynamics of EuCd_2As_2
- Summary and Acknowledgements


Fourier Transform Infrared Spectroscopy

- Source beam is split into two
 - One goes to fixed mirror; other to moving mirror
 - Switches between destructive and constructive interference
- Recombined beam is passed to sample
 - Can do transmittance (absorbance) or reflectance
- Resulting interferogram is Fourier transformed to get a spectrum in the frequency domain
- Can be used to measure over broad spectral ranges




FTIR instruments

- Different configurations for different types of measurements
 - Varying spectral ranges and resolution, spatial resolution, etc.
- Can measure different sample forms (liquid, crystalline, powders)
- Can couple in cryostats for temperature control





Raman scattering

- Similar spectral ranges but different principle than FTIR
- Excite with one laser wavelength
 - Choice can affect spectra if it is near a resonance
- Measure in relative frequency (with laser line being 0 cm^{-1})
 - Difficult to get to very low frequencies
- Can integrate over time to overcome low signal issues
 - Makes intensity relative




Measuring in extreme conditions

- Temperature (4.2 to 300 K)
 - Open-flow helium cryostat
- Magnetic field (done at NHMFL)
 - 35 T static fields
 - Up to 60 T pulsed fields
- Pressure (ambient to 20 GPa)
 - Diamond anvil cell
- Combine parameters to map out phase space






Analyzing spectra

- Fit spectra with standard functions
- Extract peak information depending on scientific interest
 - Center, fullwidth at half max, area
- Tracking as function of tuning parameter can reveal transitions
- Can be used to quantify coupling constants, fit to models, etc.





$\text{CuF}_2(\text{H}_2\text{O})_2(3\text{-chloropyridine})$

- Two-dimensional molecule-based magnet
 - Magnetic exchange within Cu-F-H₂O plane
- Magnetic crossover from AFM to FM at 0.8 GPa
- Complex spectra – many peaks in both IR and Raman




What do the high pressure spectra show?



What do we learn?


- Combined IR, Raman, and calculations to get full picture
- New hydrogen bond formed under pressure with Cl
- New magnetic exchange pathway
 - Material goes from 2D to 3D
 - Magnetism changes from AFM to FM
- New structural transition discovered near 5.5 GPa



O'Neal, Sci. Rep. **4**, 6054 (2014)


Multiferroic Ni_3TeO_6

- Ferroelectric and antiferromagnetic in ground state
- Temperature-magnetic field phase diagram known
- No previous pressure studies
- Structure is more complex than it at first appears
 - Three unique NiO_6 configurations
 - Empty O_6 cages throughout






High pressure spectra

- Measured both infrared and Raman
- Can assign all modes based on calculations and previous work




Subtle changes in spectra

- Tracking mode frequencies reveals reduced compressibility above 4 GPa
 - Slopes for several modes decrease here
- Supported by calculations of spectrum using structure under pressure
- Why only some modes?




What causes the slope change?

- Use calculated structures to look for clues
- Bond lengths all change linearly – nothing to learn
- Bond angles also change linearly – except for one!



What do we learn?



- Used detailed analysis to reveal cause of slope change
- Vibrations of atoms near the empty pocket all displayed the slope change
- Pocket compresses easily up to 4 GPa, then stiffens as atoms get closer
- Changes to magnetic exchange in line with preliminary measurements



O'Neal, Phys. Rev. B **98**, 184101 (2018)



CuGeO₃ nanorods

- Spin-Peierls transition at low temperature
- Bulk has been studied for over 20 years
- Can now grow nanorods – new way to tune the properties
 - Control of nanorod length, keeping diameter similar
 - Spin-Peierls transition suppressed in shortest rods







Vibronic coupling

- d-d electronic transition is formally forbidden
 - Activated by odd parity phonon (IR-active)
- Temperature dependent oscillator strength can be modeled
 - Can indicate which phonon(s) are activating the transition




Vibronic coupling model limitations

- Typical vibronic coupling model did not work for nanorods
- Added a temperature-independent constant, f_x
- Slight improvement in fit for bulk
 - Significant difference in phonon frequency
- Needed a physical basis for constant
 - Defects and edge effects
 - Distortions in crystal structure



What do we learn?

- New model worked for all nanorods
- Could track size dependence of coupled phonon frequency
 - Blue color comes from different origin
- Crossover in most important phonon below rod length of 500 nm
- New model can now apply to all nanomaterials




O'Neal, Phys. Rev. B 96, 075437 (2017)

Outline

- FTIR and Raman
 - Pressure-induced dimensionality crossover in Cu-3Cl(pyd)
 - Local structure changes in Ni_3TeO_6 under pressure
 - Size-dependent vibronic coupling in CuGeO_3 nanorods
- **Ultrafast spectroscopy**
 - Carrier dynamics of EuCd_2As_2
- Summary and Acknowledgements


Ultrafast spectroscopy

- Laser pulse excites material
- Second, weaker pulse measures changes
- Varying the path length of the second pulse controls its arrival time
 - Allows for ultrafast measurements even with a slow detector
- Spectral selectivity, high sensitivity, small focal size, femtosecond timescale




EuCd_2As_2 – magnetic Weyl semimetal(?)

- Magnetic derivative of Cd_2As_3
- Growth conditions affect magnetism
 - (FM) $T_c = 30$ K or (AFM) $T_N = 9$ K
- Spins forced to align out of plane leads to Weyl semimetal with one pair of nodes
- Other measurements have been inconclusive




Across the magnetic ordering transition

- Measure as function of temperature
 - Observe changes across $T_C = 30$ K
- Overall changes appear gradual
 - Magnitude rises then falls
 - Timescale follows similar trend
- Need to fit data to learn more





Fitting the data

- Fit data at each temperature to extract parameters
- Model is complex
 - Rise from laser pulsewidth
 - Slow rise process
 - Two decay processes
 - Electron-phonon relaxation
 - Spin-phonon relaxation
- Follow parameters as function of temperature to look for changes across T_C




How does magnetism affect dynamics?

- Subtle signs of spin-charge coupling across T_C
- Much stronger changes near 70 K
 - Attributed to short-range, in-plane magnetic fluctuations



Summary

- Spectroscopy is powerful and non-destructive technique that can be used to study a wide range of materials
 - Can probe phase transitions with temperature, magnetic field, and pressure
- Detailed data analysis is key to deep understanding
 - Hydrogen bond formation in $\text{CuF}_2(\text{H}_2\text{O})_2(3\text{-chloropyridine})$
 - Local structure changes in Ni_3TeO_6
 - Size-dependent vibronic coupling in CuGeO_3 nanorods
 - Ultrafast carrier dynamics in EuCd_2As_2



Acknowledgements

- $\text{CuF}_2(\text{H}_2\text{O})_2(3\text{-chloropyridine})$
 - Jan Musfeldt (advisor)
 - Puru Jena (calculations)
 - Zhenxian Liu (high pressure)
 - John Schlueter (synthesis)
- CuGeO_3
 - Jan Musfeldt (advisor)
 - Xue-Tai Chen (synthesis)
- EuCd_2As_2
 - Dmitry Yarotski (advisor)
 - Rohit Prasankumar (advisor)
 - Paul Canfield (synthesis)
- Funding
 - National Science Foundation
 - Department of Energy
 - CATS

